



| TIMI Risk Scoring                                    | Points (Circle) |
|--|-----------------|
| <b>Historical:</b>                                   |                 |
| <input type="checkbox"/> Age 65 or older             | 1               |
| <input type="checkbox"/> >3 CAD risk factors         | 1               |
| • HTN  |                 |
| • High Cholesterol                                   |                 |
| • Active Smoker                                      |                 |
| • Diabetes Mellitus                                  |                 |
| • Family History                                     |                 |
| <input type="checkbox"/> Known CAD (stenosis ≥50%)   | 1               |
| <input type="checkbox"/> ASA use in 7 days           | 1               |
| <b>Presentation:</b>                                 |                 |
| <input type="checkbox"/> Recent (<24h) severe angina | 1               |
| <input type="checkbox"/> Increase in biomarkers      | 1               |
| <input type="checkbox"/> ST deviation >0.5 mm        | 1               |
| <b>Risk Score-total points (7)</b>                   | <b>Total:</b>   |
| <b>TIMI Risk Score 0-2 : Low Risk</b>                |                 |
| <b>TIMI Risk Score 3-4 : Intermediate</b>            |                 |
| <b>TIMI Risk Score 5-7 : High Risk</b>               |                 |

**Early Management Strategy Criteria:**

|                                   |   |
|-----------------------------------|---|
| Immediate invasive (within 2 h)   | Refractory angina   |
|                                   | Signs or symptoms of HF or new or worsening mitral regurgitation                                    |
|                                   | Hemodynamic instability   |
|                                   | Recurrent angina or ischemia at rest or with low-level activities despite intensive medical therapy |
|                                   | Sustained VT or VF  |
| Ischemia-guided strategy          | Low-risk score (e.g., TIMI [0 or 1])  |
|                                   | Low-risk Tn-negative female patients  |
|                                   | Patient or clinician preference in the absence of high-risk features                                |
| Early invasive (within 24 h)      | None of the above   |
|                                   | Temporal change in Tn (Section 3.4)   |
|                                   | New or presumably new ST depression   |
| Delayed invasive (within 25-72 h) | None of the above but diabetes mellitus   |
|                                   | Renal insufficiency (GFR <60 mL/min/1.73 m <sup>2</sup> )   |
|                                   | Reduced LV systolic function (EF <0.40)   |
|                                   | Early postinfarction angina   |
|                                   | PCI within 6 mo   |
|                                   | Prior CABG  |
|                                   | TIMI score ≥2   |

D/C Patient with Cardiology follow up Appointment and Cardiac Rehab Consult